

# Response of antioxidant enzymes in *Mythimna separata* (Lepidoptera: Noctuidae) exposed to thermal stress

A. Ali<sup>1</sup>, M. A. Rashid<sup>1</sup>, Q. Y. Huang<sup>1</sup>, C. Wong<sup>2</sup> and C.-L. Lei<sup>1\*</sup>

<sup>1</sup>Hubei Insect Resources Utilization and Sustainable Pest Management Key Laboratory, College of Plant Science and Technology, Huazhong Agricultural University, Wuhan, China; <sup>2</sup>Laboratory of Pesticide Toxicology, Iowa State University, Ames, Iowa, USA

## Abstract

The oriental army worm *Mythimna separata* (Lepidoptera: Noctuidae) is a migratory pest in Eastern Asia and China. Seasonal high temperatures in Southern China and low temperatures in Northern China are pressures favouring the annual migration of this species, while cold tolerance determines the northern limit of its overwintering range. A number of physiological stress responses occur in insects as a result of variations in temperature. One reaction to thermal stress is the generation of reactive oxygen species (ROS), which can be harmful by causing oxidative damage. The time-related effects (durations of 1, 4 and 7 h) of thermal stress treatments of *M. separata* at comparatively low (5, 10, 15 and 20°C) and high (30, 35, 40 and 45°C) temperatures on the activities of antioxidant enzymes, including superoxide dismutase (SOD), catalase (CAT), peroxidase (POX) and glutathione S-transferases (GSTs), and total antioxidant capacity (T-AOC) were determined. Thermal stress resulted in significant elevation of the activities of SOD, CAT and GSTs, indicating that these enzymes contribute to defence mechanisms counteracting oxidative damage caused by an increase in ROS. However, at high-temperatures, POX and T-AOC were also found to contribute to scavenging ROS. Our results also indicate that extreme temperatures lead to elevated ROS production in *M. separata*. The present study confirms that thermal stress can be responsible for oxidative damage. To overcome such stress, antioxidant enzymes play key roles in diminishing oxidative damage in *M. separata*.

**Keywords:** thermal stress, antioxidant enzymes, oxidative stress, *Mythimna separata*

(Accepted 5 October 2016)

## Introduction

Temperature is the most critical environmental factor for many organisms; it effects growth, reproduction, distribution and abundance, by inducing numerous physiological responses (Angilletta *et al.*, 2002; Parmesan, 2006; Jia *et al.*, 2011). The thermal stress response, which occurs in all living

organisms, is a standard reaction to above normal temperatures (Kotak *et al.*, 2007; Nguyen *et al.*, 2013). Under thermal stress, overproduction of reactive oxygen species (ROS) can cause oxidative damage. In general, the production of ROS and antioxidant processes are synchronized; however, the balance between these activities can be disrupted during periods of environmental stress, leading to synthesis of additional ROS (Joanisse & Storey, 1996; Lopez-Martinez *et al.*, 2008; Lalouette *et al.*, 2011). Overproduction of ROS can disrupt the fluidity of cell membranes, due to lipid peroxidation, and lead to necrobiosis, as well as alterations in cellular DNA (Green & Reed, 1998; Monaghan *et al.*, 2009). A number of factors promote

\*Author for correspondence  
 Phone/Fax: +86 27 87287207  
 E-mail: ioir@mail.hzau.edu.cn




Fig. 1. Effects of treatment of *M. separata* adults with thermal stress for various lengths of time on CAT activity. Data collected after treatment durations of 1, 4 and 7 h are presented in (a), (b) and (c), respectively. Data are presented as means ( $\pm$ SE) of three replicate experiments. Letters above bars indicate significant differences ( $P < 0.05$ ) determined by ANOVA with Tukey's test.

overproduction of ROS in insects, including compensatory growth, ingested plant photo-oxidants and unfavourable environmental conditions (such as the presence of pollutants, adverse temperatures or hypoxic stress) (Aucoin *et al.*, 1995; Zaman *et al.*, 1995; Joannis & Storey, 1998; Jing *et al.*, 2005; Mangel & Munch, 2005).

To prevent ROS damage, living organisms have developed complex defence mechanisms for handling ROS, which include both enzymes and molecular antioxidants (Howe & Schillmiller, 2002). Anti-oxidative enzymes are the key to removal of ROS from biological systems. The primary anti-oxidative enzymes in insects are superoxide dismutase (SOD), catalase (CAT), peroxidase (POX) and glutathione-S-transferases (GSTs) (Felton & Summers, 1995; Wang *et al.*, 2001; Dubovskiy *et al.*, 2008). SOD catalyses the dismutation of superoxide radicals into oxygen and H<sub>2</sub>O<sub>2</sub>, whereas both CAT and POX catalyse the dismutation of H<sub>2</sub>O<sub>2</sub> into oxygen and water. Another important enzyme, GST, eliminates lipid peroxidation products (hydroperoxides) from cells (Dubovskiy *et al.*, 2008; Meng *et al.*, 2009). In addition, the ability of all antioxidants in an organism to counter oxidation is described as the total antioxidant capacity (T-AOC) (Ghiselli *et al.*, 2000).

The oriental army worm *Mythimna separata* (Lepidoptera: Noctuidae) is a migratory pest in Eastern Asia and China (Ruilo & Ziangshi, 1987; Rui-Lu *et al.*, 1989; Chen *et al.*,

1995). It has been responsible for damaging millet (*Pennisetum* spp.) and wheat (*Triticum* spp.) crops for thousands of years in China. Recently, it has also been found to damage rice and corn crops (Chen & Hu, 2000; Wang *et al.*, 2006). Seasonal migration of *M. separata* has been observed in China. The organism is mainly present in Southern and Central China and its population is well controlled by reducing the cultivation area of host plants in these regions, although it can survive and reproduce in some southern regions during winter. However, crops in several areas of Northern China, where the insect is unable to survive over winter, are still continuously damaged (Jiang, 2004; Zhang *et al.*, 2006). Seasonal high temperatures in Southern China and low temperatures in Northern China are one of the pressures favouring the annual migration of *M. separata* between these areas as an adaptive life history strategy (Jiang *et al.*, 2000). Conversely, the cold tolerance of this species determines the northern limit of its overwintering range in China. Zhang *et al.* (2008) revealed that cold stress (5°C) experienced during the first 24 h after eclosion can change migrant *M. separata* into resident insects. Jiang *et al.* (2011) reported that, for adults of *M. separata*, flight occurred at temperatures between 11 and 32°C, with an optimum range of 17–22°C, and a lower threshold of 8°C. Warmer temperatures generally have a positive effect on developmental




Fig. 2. Effects of treatment of *M. separata* adults with thermal stress for various lengths of time on SOD activity. Data collected after treatment durations of 1, 4 and 7 h are presented in (a), (b) and (c), respectively. Data are presented as means ( $\pm$ SE) of three replicate experiments. Letters above bars indicate significant differences ( $P < 0.05$ ) determined by ANOVA with Tukey's test.

time, lifespan, adult flight activity and reproduction of *M. separata* (Jiang & Luo, 1997; Xinfu *et al.*, 1998); however, very high temperatures can have the opposite effect, and suppress adult reproduction to a greater extent than they promote migratory flight (Jiang *et al.*, 2000). To facilitate growth and reproduction, animals search for balanced sources of nutrition, mates and oviposition sites. This kind of searching behaviour has costs that are offset by the benefits gained from the resource (Crespo *et al.*, 2014). *M. separata* encounters thermal fluctuations during its life cycle. The cost to the adults of extreme temperatures (both low and high) is much higher than that of migratory flight. To date, the effects of thermal stress on *M. separata* have not been reported. The aim of the present study was to determine how variations in temperature affect anti-oxidant enzyme activities in response to oxidative stress as such changes may lead *M. separata* to migrate in order to survive in different seasons.

## Materials and methods

### Insects

Insects for experimentation were collected from the Key Laboratory of Insect Resources Utilization and

Sustainable Pest Management, Huazhong Agricultural University, Wuhan. *M. separata* were reared at room temperature ( $25 \pm 2^\circ\text{C}$ ),  $60 \pm 10\%$  relative humidity, and with 14:10 h light:dark cycles. An artificial diet was used to feed the larvae as described in Chun (1981).

### Thermal stress

Three-day-old adults were selected for the experiment. Five adults were transferred into 100 ml plastic containers for each treatment. Insects underwent temperature treatments, at 5, 10, 15, 20, 30, 35, 40 and  $45^\circ\text{C}$ , for 1, 4 and 7 h. For all stress treatments, a programmable thermal controller (Ningbo Southeast Instrument, RXZ-260B, China) was used. A temperature of  $25^\circ\text{C}$  was set as the control for this experiment. Adult insects were frozen in liquid nitrogen immediately after temperature treatment and stored at  $-80^\circ\text{C}$  until further analysis. Experiments were performed three times on three different days.

### Enzyme extraction

A commercially available assay kit (Nanjing Jiancheng Bioengineering Institute, China) was used for extraction of enzymes, according to the manufacturer's instructions. Samples




Fig. 3. Effects of treatment of *M. separata* adults with thermal stress for various lengths of time on POX activity. Data collected after treatment durations of 1, 4 and 7 h are presented in (a), (b) and (c), respectively. Data are presented as means ( $\pm$ SE) of three replicate experiments. Letters above bars indicate significant differences ( $P < 0.05$ ) determined by ANOVA with Tukey's test.

were homogenized in 0.9% saline solution at a ratio of 1:9 ( $W_{\text{flies}}:V_{\text{normal saline}}$ ). Homogenates were centrifuged at 10,000  $g$  for 15 min at 4°C. After centrifugation, the supernatant was stored at low temperature until tested to determine enzyme activity. The method of Bradford (1976) was used to calculate protein concentrations.

#### Measurement of T-AOC

T-AOC was measured using an assay kit (Nanjing Jiancheng Bioengineering Institute) according to the manufacturer's instructions. The kit is based on the ability of antioxidant substances present in the supernatant to reduce a pool of ferric iron. This acts as a redox-linked, reductant colorimetric assay, as a relatively stable complex is formed between  $\text{Fe}^{2+}$  and porphyrin, which absorbs light at 520 nm. The required quantity of protein to elevate the absorbance measurement by 0.01 nm  $\text{min}^{-1}$  mg<sup>-1</sup> protein was defined as one unit of T-AOC.

#### Determination of antioxidant enzyme activities

Spectrophotometry was used to determine the activities of enzymes (SOD, CAT, POX and GST) using assay kits (Nanjing

Jiancheng Bioengineering Institute), in accordance with the instructions of the manufacturer.

CAT activity was calculated by gauging the decline in absorbance at 405 nm in response to decomposition of  $\text{H}_2\text{O}_2$ . The amount of enzyme required for decomposition of  $\text{H}_2\text{O}_2$  per second per mg of protein was defined as one unit of CAT activity. The unit of expression for CAT activity was U mg<sup>-1</sup> protein.

The xanthine oxidase method was used to determine SOD activity at 450 nm. The quantity of enzyme required for 50% inhibition of the xanthine-xanthine oxidase reaction in a protein concentration of 1 mg ml<sup>-1</sup> was defined as one unit of SOD activity, expressed as U mg<sup>-1</sup> protein.

POX activity was measured at 420 nm by the activation of oxidation in the presence of  $\text{H}_2\text{O}_2$ . The quantity of POX enzyme required to catalyse 1  $\mu\text{g}$  substrate  $\text{min}^{-1}$  mg<sup>-1</sup> of protein was defined as one unit of POX activity, and expressed as U mg<sup>-1</sup> protein.

The substrate, 1-chloro-2,4-dinitrobenzene (CDNB) was used to determine the activity of GST. A change in absorbance at 412 nm was observed due to the formation of GSH-CDNB. The amount of GST enzyme required to activate the fusion of 1  $\mu\text{mol l}^{-1}$  GSH with CDNB  $\text{min}^{-1}$  mg<sup>-1</sup> protein was




Fig. 4. Effects of treatment of *M. separata* adults with thermal stress for various lengths of time on GST activity. Data collected after treatment durations of 1, 4 and 7 h are presented in (a), (b) and (c), respectively. Data are presented as means ( $\pm$ SE) of three replicate experiments. Letters above bars indicate significant differences ( $P < 0.05$ ) determined by ANOVA with Tukey's test.

defined as one unit of GST activity and expressed as  $\text{U mg}^{-1}$  protein.

#### Statistical analysis

Treatment effects (temperature and duration) were subjected to one or two-way analysis of variance (ANOVA) using the general linear model procedure in SPSS 16.0 (SPSS, Chicago, IL, USA); when significant effects were identified, mean differences were separated by Tukey's test, with  $P < 0.05$  considered statistically significant.

### Results

At 45°C, all adults died, regardless of the duration of the thermal stress treatment.

#### Antioxidant enzymes

CAT activity in *M. separata* adults was significantly increased at both low and high, compared with the control, temperatures ( $P < 0.01$ ), after treatment for all durations ( $P < 0.01$ ), and the interaction between temperature and duration was significant ( $P < 0.01$ ). Maximum CAT activity values were 167.67, 146.94 and 135.50  $\text{U mg}^{-1}$  protein recorded under cold stress (5°C) for 1, 4 and 7 h, respectively (fig 1).

SOD activity was significantly raised at both low and high, compared with the control, temperatures in *M. separata* adults ( $P < 0.01$ ), for all durations of treatment ( $P < 0.01$ ), and there was a significant interaction between temperature and duration ( $P < 0.01$ ). The highest SOD activity levels (53.92, 67.41 and 69.45  $\text{U mg}^{-1}$  protein) were observed under cold stress (5°C) for 1, 4 and 7 h, respectively (fig 2).

POX activity in *M. separata* adults was also significantly affected at all temperatures ( $P < 0.01$ ) and for all durations ( $P < 0.01$ ), and temperature and duration interacted significantly ( $P < 0.01$ ). POX activity increased significantly under high-temperature stress (temperatures ranging from 30 to 40°C) at 1 and 4 h, relative to cold stress and control (25°C) conditions; however, after 7 h, while a significant elevation in POX activity was observed at temperatures of 15, 20, 30, 35 and 40°C, relative to that at 10°C, no significant differences were observed at either low or high temperatures compared with the control group (25°C) (fig 3).

Significant increases in GST activity in *M. separata* adults at both low and high temperatures were observed at all temperatures ( $P < 0.01$ ) and durations ( $P < 0.01$ ), compared with controls (25°C), and there was a significant interaction between temperature and durations ( $P < 0.01$ ). The highest values of GST activity recorded were 649.71 and 572.50  $\text{U mg}^{-1}$  protein at 5°C for 1 and 4 h, respectively. In addition, after 7 h at 30°C GST activity was 633.15  $\text{U mg}^{-1}$  protein (fig 4).




Fig. 5. Effects of treatment of *M. separata* adults with thermal stress for various lengths of time on total antioxidant capacity. Data collected after treatment durations of 1, 4 and 7 h are presented in (a), (b) and (c), respectively. Data are presented as means ( $\pm$ SE) of three replicate experiments. Letters above bars indicate significant differences ( $P < 0.05$ ) determined by ANOVA with Tukey's test.

#### Total antioxidant capacity (T-AOC)

Significant effects were observed on the T-AOC of *M. separata* adults, relative to the control group, under both low- and high-temperature stresses ( $P < 0.01$ ). The duration of treatment did not result in a significant change in T-AOC ( $P < 0.31$ ); however, a significant interaction between temperature and duration was observed ( $P < 0.01$ ). Heat stress (temperatures ranging from 30 to 40°C) resulted in a significant increase in T-AOC after 1 h, relative to cold stress and control temperature; however, after 4 h of treatment only temperature stress treatment at 40°C resulted in significantly increased T-AOC compared with controls (25°C). No significant changes were observed compared with controls when adults were exposed to low- and high-temperature stresses for 7 h (fig 5).

#### Discussion

Temperature is a critical environmental variable that engenders physiological changes in organisms (Jia *et al.*, 2011). *M. separata* adults were exposed to different thermal stresses, at both low and high temperatures, and consequent physiological oxidative stress responses explored. The effect of different thermal stress conditions on the activities of the antioxidant enzymes, SOD, CAT, POX, GST and on T-AOC, in *M. separata* adults was examined. CAT, SOD, POX and

GST are key antioxidant defence enzymes, which work in a synchronized manner to thwart oxidative stress caused by high concentrations of ROS within cells. Among these antioxidant enzymes, CAT is considered to be the principle H<sub>2</sub>O<sub>2</sub> scavenging enzyme in arthropods (Jena *et al.*, 2013), as selenium-dependent glutathione POX (the main catalyser in other organisms) is deficient (Sohal *et al.*, 1990). However, CAT is ineffective for the removal of low concentrations of H<sub>2</sub>O<sub>2</sub>, as it functions only in the presence of high cellular concentrations (Ahmad *et al.*, 1991). Under thermal stress, CAT activity in citrus red mites is insufficient (Yang *et al.*, 2010); however, in the present study, a significant elevation of CAT activity was observed at both low and high temperatures in *M. separata* adults, compared with controls. These data suggest that overexpression of CAT enhances the removal of H<sub>2</sub>O<sub>2</sub> at both low and high temperatures, and prevents oxidative stress damage. Similar results were reported by Jia *et al.* (2011), and Nabizadeh & Kumar (2011), in the oriental fruit fly, *Bactrocera dorsalis* and the silkworm, *Bombyx mori*.

SOD plays a critical role in reducing high levels of superoxide radicals induced by exposure to low and high temperatures (Celino *et al.*, 2011). In the present study, significant enhancement of SOD activity was determined under conditions of thermal stress, compared with controls at 25°C, suggesting that SOD production was induced as a result of temperature fluctuations to protect *M. separata* adults from

thermal stress. Similar results were reported by McCord & Fridovich (1969) and Jia *et al.* (2011). SOD and CAT can directly remove excess ROS in a coordinated manner. SOD removes O<sub>2</sub><sup>-</sup> through the process of dismutation to O<sub>2</sub> and H<sub>2</sub>O<sub>2</sub>, and H<sub>2</sub>O<sub>2</sub> is then sequentially reduced to H<sub>2</sub>O and O<sub>2</sub> by CAT (Kashiwagi *et al.*, 1997). The observed higher levels of CAT, relative to those of SOD, in this study indicate that, under thermal stress, H<sub>2</sub>O<sub>2</sub> is also synthesized by processes other than SOD activity.

GSTs can metabolize lipid peroxidation products together with POX, which also breaks down H<sub>2</sub>O<sub>2</sub> (Jia *et al.*, 2011). In the present study, POX activity increased significantly at high temperatures (ranging from 30 to 40°C) for 1 and 4 h, compared with controls. Similar findings were reported by Zhang *et al.* (2014) in the predatory mite, *Neoseiulus cucumeris*. Our results demonstrate that POX activity was expeditiously induced by thermal stress in *M. separata* adults, which is consistent with the findings of a similar study involving *Helicoverpa armigera* (Meng *et al.*, 2009). However, after the longest duration (7 h) of thermal stress, a significant decrease in POX activity was observed in the oriental fruit fly *B. dorsalis* (Jia *et al.*, 2011) and predatory mite, *N. cucumeris* (Zhang *et al.*, 2014). In contrast, our results indicate no significant changes in POX activity at either low or high temperatures compared with the control temperature after the longest treatment duration (7 h), similar to the results reported by Yang *et al.* (2010). The elevation of POX activity at higher temperatures indicates that it was stimulated by scavenging ROS in *M. separata*.

GSTs are a group of multifunctional dimeric enzymes, which catalyse the conjugation of glutathione to a broad spectrum of endogenous and xenobiotic compounds for detoxification, protection from oxidative damage, isomerization and intercellular transportation (Board & Menon, 2013). These enzymes are involved in the inactivation of toxic lipid peroxidation products created by oxidative stress damage. In the present study, the observation of significantly elevated levels of GST under temperature stress suggests that this enzyme protects *M. separata* adults from oxidative damage under these conditions. Similar antioxidant responses have been reported in *P. japonica* (Zhang *et al.*, 2015), *A. mylitta* (Jena *et al.*, 2013), *B. dorsalis* (Jia *et al.*, 2011) and *P. citri* (Yang *et al.*, 2010).

T-AOC is widely used as a tool to assess redox, and as a representative measure of the total antioxidant capacity existing in an organism (Meng *et al.*, 2009; Yang *et al.*, 2010; Sashidhara *et al.*, 2011). T-AOC was augmented significantly when *M. separata* adults were exposed to high temperatures (ranging from 30 to 40°C) for 1 h and (40°C) for 4 h, compared with controls. These data suggest that T-AOC adapts to deal with oxidative stress and free radical formation and are consistent with the results reported by Zhang *et al.* (2015), Zhang *et al.* (2014) and Jia *et al.* (2011). However, no significant difference was observed compared with controls after treatment for the longest duration (7 h). A similar result was reported by Jia *et al.* (2011) in *B. dorsalis* under thermal stress conditions.

Antioxidant stress is well managed by antioxidant enzymes; however, some non-enzymatic substances, e.g. trehalose (Mahmud *et al.*, 2010) and vitamin E (α-tocopherol) (Kaur *et al.*, 2009) also contribute to this process. A recent study also confirmed the involvement of heat shock proteins, along with antioxidant enzymes, in the response to ROS damage (Rosa *et al.*, 2012). The increase of T-AOC only at high temperatures indicates that *M. separata* uses not only antioxidant enzymes,

but also other defence mechanisms, to combat thermal stress and enable survival of the organism (Jia *et al.*, 2011).

## Conclusion

Oxidative stress can be generated when environmental factors disturb the balance of redox reactions within an organism. In *M. separata*, thermal stress is the main candidate factor for the induction of oxidative stress. In response to thermal stress, antioxidant enzymes are upregulated as a defence mechanism to mitigate potential cellular damage. The enzymes SOD, CAT and GST undergo significant increases in activity in response to thermal stress in *M. separata*, and may be involved in the management of oxidative damage produced by ROS. Indeed, there was an increased production of ROS at higher temperatures; therefore, these fluctuations may reflect physiological adaptations in *M. separata* related to its migration habits. However, at high temperatures, compared with lower temperatures, POX activity and T-AOC have additional roles in scavenging ROS.

## Acknowledgements

This study was supported by the National Department Public Benefit (Agriculture) Research Foundation (Grant No. 201403031) and the National Natural Science Foundation of China (Grant No. 31572017).

## References

- Ahmad, S., Duval, D.L., Weinhold, L.C. & Pardini, R.S. (1991) Cabbage looper antioxidant enzymes: tissue specificity. *Insect Biochemistry* **21**, 563–572.
- Angilletta, M.J., Niewiarowski, P.H. & Navas, C.A. (2002) The evolution of thermal physiology in ectotherms. *Journal of Thermal Biology* **27**, 249–268.
- Aucoin, R., Guillet, G., Murray, C., Philogène, B.J. & Arnason, J. T. (1995) How do insect herbivores cope with the extreme oxidative stress of phototoxic host plants. *Archives of Insect Biochemistry and Physiology* **29**, 211–226.
- Board, P.G. & Menon, D. (2013) Glutathione transferases, regulators of cellular metabolism and physiology. *Biochimica et Biophysica Acta* **1830**, 3267–3288.
- Bradford, M.M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* **72**, 248–254.
- Celino, F.T., Yamaguchi, S., Miura, C., Ohta, T., Tozawa, Y., Iwai, T. & Miura, T. (2011) Tolerance of spermatogonia to oxidative stress is due to high levels of Zn and Cu/Zn superoxide dismutase. *PLoS ONE* **6**, e16938.
- Chen, S.D. & Hu, B.H. (2000) *Plant Protection in China in Fifty Years*. Beijing, China, Agriculture Press.
- Chen, R., Sun, Y., Wang, S., Zhai, B. & Bao, X. (1995) *Mythimna Separata in East Asia in Relation to Weather and Climate. I. Northeastern China. Insect Migration: Tracking Resources through Space and Time*. Cambridge, Cambridge University Press, pp. 93–104.
- Chun, B.F. (1981) A new artificial diet for army worm. *Acta Entomologica Sinica* **24**, 379–338.
- Crespo, J.G., Vickers, N.J. & Goller, F. (2014) Male moths optimally balance take-off thoracic temperature and warm-up duration to reach a pheromone source quickly. *Animal Behaviour* **98**, 79–85.

- Dubovskiy, I., Martemyanov, V., Vorontsova, Y., Rantala, M., Gryzanova, E. & Glupov, V. (2008) Effect of bacterial infection on antioxidant activity and lipid peroxidation in the midgut of *Galleria mellonella* L. larvae (Lepidoptera, Pyralidae). *Comparative Biochemistry and Physiology C: Toxicology & Pharmacology* **148**, 1–5.
- Felton, G.W. & Summers, C.B. (1995) Antioxidant systems in insects. *Archives of Insect Biochemistry and Physiology* **29**, 187–197.
- Ghiselli, A., Serafini, M., Natella, F. & Scaccini, C. (2000) Total antioxidant capacity as a tool to assess redox status: critical view and experimental data. *Free Radical Biology and Medicine* **29**, 1106–1114.
- Green, D.R. & Reed, J.C. (1998) Mitochondria and apoptosis. *Science* **281**, 1309.
- Howe, G.A. & Schillmiller, A.L. (2002) Oxylin metabolism in response to stress. *Current Opinion in Plant Biology* **5**, 230–236.
- Jena, K., Kar, P. K., Kausar, Z. & Babu, C.S. (2013) Effects of temperature on modulation of oxidative stress and antioxidant defenses in testes of tropical tasar silkworm *Antheraea mylitta*. *Journal of Thermal Biology* **38**, 199–204.
- Jia, F.X., Dou, W., Hu, F. & Wang, J.J. (2011) Effects of thermal stress on lipid peroxidation and antioxidant enzyme activities of oriental fruit fly, *Bactrocera dorsalis* (Diptera: Tephritidae). *Florida Entomologist* **94**, 956–963.
- Jiang, X. (2004) The physiological and genetic characteristics of migratory behavior and genetic diversity, as determined by AFLP in the oriental armyworm, *Mythimna separata* (Walker). PhD dissertation, Chinese Academy of Agricultural Sciences, Beijing, China.
- Jiang, X.F. & Luo, L.Z. (1997) Influence of temperature at pupal and adult stage on flight capacity of adult oriental armyworm, *Mythimna separata* (Walker). pp. 274–280 in Chen, X.F., Dai, X.F. & Hu, T. (Eds) *Ecological Research Sustainable Development*. Beijing, China, China Environmental Science Press.
- Jiang, X.F., Luo, L.Z. & Hu, Y. (2000) Influences of rearing temperature on flight and reproductive capacity of adult oriental armyworm, *Mythimna separata* (Walker). *Acta Ecologica Sinica* **20**, 288–292.
- Jiang, X., Luo, L., Zhang, L., Sappington, T.W. & Hu, Y. (2011) Regulation of migration in *Mythimna separata* (Walker) in China: a review integrating environmental, physiological, hormonal, genetic, and molecular factors. *Environmental Entomology* **40**, 516–533.
- Jing, X.H., Wang, X.H. & Kang, L. (2005) Chill injury in the eggs of the migratory locust, *Locusta migratoria* (Orthoptera: Acrididae): the time temperature relationship with high temperature interruption. *Insect Science* **12**, 171–178.
- Joanisse, D. & Storey, K. (1996) Oxidative stress and antioxidants in overwintering larvae of cold-hardy goldenrod gall insects. *Journal of Experimental Biology* **199**, 1483–1491.
- Joanisse, D.R. & Storey, K.B. (1998) Oxidative stress and antioxidants in stress and recovery of cold-hardy insects. *Insect Biochemistry and Molecular Biology* **28**, 23–30.
- Kashiwagi, A., Kashiwagi, K., Takase, M., Hanada, H. & Nakamura, M. (1997) Comparison of catalase in diploid and haploid *Rana rugosa* using heat and chemical inactivation techniques. *Comparative Biochemistry and Physiology B: Biochemistry and Molecular Biology* **118**, 499–503.
- Kaur, G., Alam, M.S. & Athar, M. (2009) Cumene hydroperoxide debilitates macrophage physiology by inducing oxidative stress: possible protection by  $\alpha$ -tocopherol. *Chemico-Biological Interactions* **179**, 94–102.
- Kotak, S., Larkindale, J., Lee, U., Von Koskull-Döring, P., Vierling, E. & Scharf, K.D. (2007) Complexity of the heat stress response in plants. *Current Opinion in Plant Biology* **10**, 310–316.
- Lalouette, L., Williams, C., Hervant, F., Sinclair, B.J. & Renault, D. (2011) Metabolic rate and oxidative stress in insects exposed to low temperature thermal fluctuations. *Comparative Biochemistry and Physiology A: Molecular & Integrative Physiology* **158**, 229–234.
- Lopez-Martinez, G., Elnitsky, M.A., Benoit, J.B., Lee, R.E. & Denlinger, D.L. (2008) High resistance to oxidative damage in the Antarctic midge *Belgica antarctica*, and developmentally linked expression of genes encoding superoxide dismutase, catalase and heat shock proteins. *Insect Biochemistry and Molecular Biology* **38**, 796–804.
- Mahmud, S.A., Hirasawa, T. & Shimizu, H. (2010) Differential importance of trehalose accumulation in *Saccharomyces cerevisiae* in response to various environmental stresses. *Journal of Bioscience and Bioengineering* **109**, 262–266.
- Mangel, M. & Munch, S.B. (2005) A life history perspective on short and long term consequences of compensatory growth. *The American Naturalist* **166**, E155–E176.
- McCord, J.M. & Fridovich, I. (1969) Superoxide dismutase an enzymic function for erythrocyte (hemocuprein). *Journal of Biological Chemistry* **244**, 6049–6055.
- Meng, J.Y., Zhang, C.Y., Zhu, F., Wang, X.P. & Lei, C.L. (2009) Ultraviolet light-induced oxidative stress: effects on antioxidant response of *Helicoverpa armigera* adults. *Journal of Insect Physiology* **55**, 588–592.
- Monaghan, P., Metcalfe, N.B. & Torres, R. (2009) Oxidative stress as a mediator of life history trade-offs: mechanisms, measurements and interpretation. *Ecology Letters* **12**, 75–92.
- Nabizadeh, P. & Kumar, T.J. (2011) Fat body catalase activity as a biochemical index for the recognition of thermotolerant breeds of mulberry silkworm, *Bombyx mori* L. *Journal of Thermal Biology* **36**, 1–6.
- Nguyen, T.M., Bressac, C. & Chevrier, C. (2013) Heat stress affects male reproduction in a parasitoid wasp. *Journal of Insect Physiology* **59**, 248–254.
- Parmesan, C. (2006) Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology, Evolution, and Systematics* **37**, 637–669.
- Rosa, R., Pimentel, M.S., Boavida, P.J., Teixeira, T., Trübenbach, K. & Diniz, M. (2012) Ocean warming enhances malformations, premature hatching, metabolic suppression and oxidative stress in the early life stages of a keystone squid. *PLoS ONE* **7**, e38282.
- Ruilo, C. & Ziangshi, B. (1987) Research on the migration of oriental armyworm in China and a discussion of management strategy. *International Journal of Tropical Insect Science* **8**, 571–572.
- Rui-Lu, C., Xiang-zhe, B., Drake, V., Farrow, R., Su-Yun, W., Ya-Jie, S. & Bao-Ping, Z. (1989) Radar observations of the spring migration into northeastern China of the oriental armyworm moth, *Mythimna separata*, and other insects. *Ecological Entomology* **14**, 149–162.
- Sashidhara, K.V., Singh, S.P., Srivastava, A. & Puri, A. (2011) Identification of the antioxidant principles of *Polyalthia longifolia* var. pendula using TEAC assay. *Natural Product Research* **25**, 918–926.
- Sohal, R., Arnold, L. & Orr, W.C. (1990) Effect of age on superoxide dismutase, catalase, glutathione reductase, inorganic peroxides, TBA-reactive material, GSH/GSSG, NADPH/

- NADP<sup>+</sup> and NADH/NAD<sup>+</sup> in *Drosophila melanogaster*. *Mechanisms of Ageing and Development* **56**, 223–235.
- Wang, Y., Oberley, L.W. & Murhammer, D.W.** (2001) Antioxidant defense systems of two lepidopteran insect cell lines. *Free Radical Biology and Medicine* **30**, 1254–1262.
- Wang, G.P., Zhang, Q.W., Ye, Z.H. & Luo, L.Z.** (2006) The role of nectar plants in severe outbreaks of armyworm *Mythimna separata* (Lepidoptera: Noctuidae) in China. *Bulletin of Entomological Research* **96**, 445–455.
- Xinfu, J., Yueqiu, L. & Lizhi, L.** (1998) Effects of high temperature on the immature stages of the oriental armyworm *Mythimna separata* Walker. *Journal of Beijing Agricultural College (China)* **13**, 20–26.
- Yang, L.H., Huang, H. & Wang, J.J.** (2010) Antioxidant responses of citrus red mite, *Panonychus citri* (McGregor) (Acari: Tetranychidae), exposed to thermal stress. *Journal of Insect Physiology* **56**, 1871–1876.
- Zaman, K., MacGill, R.S., Johnson, J.E., Ahmad, S. & Pardini, R.S.** (1995) An insect model for assessing oxidative stress related to arsenic toxicity. *Archives of Insect Biochemistry and Physiology* **29**, 199–209.
- Zhang, L., Luo, L.Z., Jiang, X.F. & Hu, Y.** (2006) Influences of starvation on the first day after emergence on ovarian development and flight potential in adults of the oriental armyworm, *Mythimna separata* (Walker) (Lepidoptera: Noctuidae). *Acta Entomologica Sinica* **49**, 895–902.
- Zhang, L., Jiang, X.F. & Luo, L.Z.** (2008) Determination of sensitive stage for switching migrant armyworms into residents. *Environmental Entomology* **37**, 1389–1395.
- Zhang, G.H., Liu, H., Wang, J.J. & Wang, Z.Y.** (2014) Effects of thermal stress on lipid peroxidation and antioxidant enzyme activities of the predatory mite, *Neoseiulus cucumeris* (Acari: Phytoseiidae). *Experimental and Applied Acarology* **64**, 73–85.
- Zhang, S., Fu, W., Li, N., Zhang, F. & Liu, T.X.** (2015) Antioxidant responses of *Propylaea japonica* (Coleoptera: Coccinellidae) exposed to high temperature stress. *Journal of Insect Physiology* **73**, 47–52.